



**Europäisches
Patentamt**

**European
Patent Office**

PCT/EP 03/34948
Rec'd PCT/PTO 06 MAY 2005 #2

**Office européen
des brevets**

REC'D 22 DEC 2003

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02079648.8

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:
Application no.: 02079648.8
Demande no:

Anmeldetag:
Date of filing: 07.11.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Akzo Nobel N.V.
Velperweg 76
6824 BM Arnhem
PAYS-BAS

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Non-steroidal androgen compounds

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C07D209/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

NON-STEROIDAL ANDROGEN COMPOUNDS

The invention relates to indole derivatives, their preparation and their use for the treatment of androgen-receptor related conditions, disorders or diseases.

5

Compounds possessing androgenic activity are useful in the treatment of men with low endogenous levels of circulating androgenic hormones or men with suppressed androgenic effects. Such treatments are prescribed to older men, to hypogonadal men or men treated with progestagens for male contraception. In addition, potent androgens suppress spermatogenesis and can be used as male contraceptives.


10

It is thus important to obtain compounds with high affinity for the androgen receptor. Non-steroidal compounds with high affinity for the androgen receptor are particularly useful since they have different tissue distribution characteristics than steroidal androgens and can be designed by proper choice of substituents to be more or less selective for certain tissues. For example, an action in the brain is usually prevented when compounds are strongly hydrophilic or carry an ionic charge.

15

The subject invention provides non-steroidal compounds with affinity for the androgen receptor. These compounds are useful for the treatment of androgen-receptor related disorders. The compounds of the subject invention have a structure according to formula I:

20



wherein

25

X is S, SO or SO₂;

R¹ is a 5-6-membered monocyclic, hetero- or homocyclic, saturated or unsaturated ring structure optionally substituted with one or more substituents selected from

the group consisting of halogen, CN, (1C-4C)fluoroalkyl, nitro, (1C-4C)alkyl, (1C-4C)alkyloxy or (1C-4C)fluoroalkoxy;

R^2 is 2-nitrophenyl, 2-cyanophenyl, 2-hydroxymethyl-phenyl, 2-pyridyl, 2-pyridyl-N-oxide, 2-benzamide, 2-benzoic acid methyl ester or 2-methoxyphenyl;

5 R^3 is H, halogen or (1C-4C)alkyl;

R^4 is H, OH or halogen;

R^5 is H, OH, (1C-4C)alkoxy, NH_2 , CN, halogen, (1C-4C)fluoroalkyl, NO_2 , hydroxy(1C-4C)alkyl, CO_2H , NHR^6 or $C(O)N(R^8, R^9)$. Herein is R^6 (1C-6C)acyl optionally substituted with one or more halogens, and are R^8 and R^9 each
10 independently H, (1C-6C)cycloalkyl, CH_2R^{10} or form together with the N a heterocyclic 5-6-membered saturated or unsaturated ring optionally substituted with (1C-4C)alkyl, where by R^{10} is H, (1C-5C)alkyl, hydroxy(1C-3C)alkyl, (1C-4C)alkylester of carboxy(1C-4C)alkyl, (1C-3C)alkoxy(1C-3C)alkyl, (mono- or di(1C-4C)alkyl)amino, (mono- or di(1C-4C)alkyl)aminocarbonyl, or a 5-6-
15 membered monocyclic, homo- or heterocyclic, aromatic or non-aromatic ring.

In one embodiment R^1 is a 5-6-membered monocyclic, hetero- or homocyclic, saturated or unsaturated ring structure optionally substituted with one or more substituents selected from the group consisting of halogen, CN, CF_3 , nitro, methoxy,
20 trifluoromethoxy or methyl; and R^2 is 2-nitrophenyl, 2-cyanophenyl, 2-hydroxymethyl-phenyl, 2-pyridyl, 2-pyridyl-N-oxide, benzamide, 2-benzoic acid methyl ester or 2-methoxyphenyl; and R^3 is H, halogen or (1C-2C)alkyl; and R^4 is H or F.

In another embodiment R^5 is H, (1C-4C)alkoxy, CN, Hal, (1C-4C)fluoroalkyl, NO_2 , or
25 R^5 is NHR^6 , wherein R^6 is (1C-6C)acyl optionally substituted with one or more halogens, or R^5 is $C(O)N(R^8, R^9)$, wherein R^8 and R^9 each independently are H, (1C-6C)cycloalkyl, or R^8 and R^9 form together with the N a heterocyclic 5-6-membered saturated or unsaturated ring optionally substituted with (1C-4C)alkyl, or R^8 and R^9
30 each independently are CH_2R^{10} , wherein R^{10} is H, (1C-5C)alkyl, hydroxy(1C-3C)alkyl, carboxy(1C-4C)alkyl, (1C-3C)alkoxy(1C-3C)alkyl, (mono- or di(1C-4C)alkyl)amino, (mono- or di(1C-4C)alkyl)-aminocarbonyl, or a 5-6-membered monocyclic, homo- or heterocyclic, aromatic or non-aromatic ring.

In yet another embodiment R^3 is H or halogen; R^4 being H; and R^5 being H, OH, (1C-4C)alkyloxy, CN, F, Cl, CF_3 , NO_2 , or R^5 is NHR^6 , wherein R^6 is (1C-3C)acyl optionally substituted with one or more halogens, or R^5 is $C(O)N(R^8, R^9)$, wherein R^8 and R^9 each
35

The androgen receptor affinity and efficacy of the compounds according to the invention makes them suitable for use in the treatment of androgen-receptor related disorders and in diagnostic methods focussed on the amount and/or location of androgen receptors in various tissues. For the latter purpose it can be preferred to make labelled variants of the compounds according to the invention. Typical androgen receptor-related treatments are those for male contraception and male or female hormone replacement therapy. Thus the invention also pertains to a method of treatment of androgen insufficiency, by administering to a male or female human or animal an effective amount of any of the above compounds. The subject invention also lies in the use of any of its compounds for the preparation of a medicine for treating androgen insufficiency. In the context of the invention, the term "androgen insufficiency" is to be understood to pertain to all kinds of diseases, disorders, and symptoms in which a male or a female suffers from too low a testosterone level, such as in hypogonadal men. In particular, the androgen insufficiency to be treated by a compound of the invention is the reduction of the testosterone level which a male individual incurs as a result of age (the compound of the invention is then used for male hormone replacement therapy), or when he is subject to male contraception. In the context of male contraception, the compound of the invention especially serves to neutralise the effect of regimens of male hormone contraception in which a sterilant such as a progestagen or LHRH (luteinizing hormone releasing hormone) is administered regularly, e.g. daily, or it is used as the sole male contraceptive substance.

The compounds of the invention can be produced by various methods known in the art of organic chemistry in general. More specifically the routes of synthesis as illustrated in the following schemes and examples can be used. In the schemes and examples the following abbreviations were used:

DMF = dimethylformamide

mCPBA = *meta* chloro perbenzoic acid

THF = tetrahydrofuran

TBTU = 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate

DIPEA = diisopropylethylamine

EDCI = 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride

HOBt = 1-hydroxybenzotriazole

NMM = N-methylmorpholine

SPE = solid phase extraction

RP-SPE = reversed phase solid phase extraction

DMSO = dimethylsulfoxide

DCM = dichloromethane


DHT = 5 α -dihydrotestosterone

5


In each of the Schemes I-V the meanings of the symbols correspond to the definitions given in the previous paragraphs.

10 Substituted indole compounds of structure 3 were prepared in two steps from the correctly substituted indoles of structure 1, via two different routes. In the first route a correctly substituted indole of structure 1 is N-alkylated with a halide of type R^1CH_2Y , where Y is a halogen, with NaH as a base in DMF at 0°C to room temperature, to give a compound of structure 2. Structure 2 is then sulfonylated at the C-3 position of the indole ring by reaction with a sulfonyl chloride in either CH_2Cl_2 or diethyl ether at room
15 temperature, to give structure 3. In the second route the two steps of the first route are reversed: the correctly substituted indole of structure 1 is first sulfonylated at the C-3 position of the indole to give structure 4, followed by N-alkylation with R^1CH_2Y to give structure 3 (Scheme I).

Scheme I




Sulfoxides of structure 5 can be obtained by oxidation of the corresponding sulfide (Structure 3) by reaction with 0.9 equivalents of mCPBA. Sulfones of structure 6 can be obtained by oxidation of the corresponding sulfide (Structure 3) by reaction with 3 equivalents of mCPBA (Scheme II).



Scheme II

10 Scheme III describes the synthesis of compounds of structure 3, in which the R² group is a phenyl ring substituted on the 2-position with either CO₂Me (Structure 7), CH₂OH (Structure 8), CO₂H (Structure 9), CONHMe (Structure 10), CONH₂ (Structure 11) or CN (Structure 12).




Scheme III

- 5 In the first step a substituted indole of structure 6 is sulfenylated at the 3-position with 2-(carboxymethyl)-phenylsulfenyl chloride, which was prepared from methylthiosalicylate and chlorine gas, to give a compound of structure 7. Reduction of the methyl ester moiety of compounds of structure 7 with LiAlH_4 gave the corresponding hydroxymethyl compounds of structure 8. Saponification of the methyl ester moiety of compounds of structure 7 with lithium hydroxide gave the corresponding carboxylic acid compounds of structure 9. The carboxylic acid moiety of
- 10


compounds of structure 9 can be converted to the corresponding carboxamide by reaction with an amine, TBTU and DIPEA in DMF at room temperature. By this method structure 10 was obtained when methylamine was used as the amine and structure 11 was obtained when NH_3 was used as the amine. Dehydration of the benzamide moiety of structure 11 with TF_2O and triethylamine in CH_2Cl_2 afforded the corresponding benzonitrile compound of structure 12 (Scheme III).

Scheme IV



Scheme IV describes the synthesis of compounds of structure 3, in which the indole ring is substituted at the 6-position with either CO_2Me (Structure 13), CO_2H (Structure 14), CONR^8R^9 (Structure 15), CONH_2 (Structure 16) or CN (Structure 17).

Saponification of the methyl ester moiety of structure 13 with lithium hydroxide afforded the corresponding carboxylic acid of structure 14. The carboxylic acid moiety of structure 14 can be converted to the corresponding carboxamides of structures 15 and 16 by reaction with an amine or amine salt, in the presence of EDCI and HOBT in a mixture of NMM and DMF at room temperature. Dehydration of the benzamide moiety of structure 16 with TF_2O and triethylamine in CH_2Cl_2 afforded the corresponding benzonitrile compound of structure 17 (Scheme IV).



Scheme V describes the synthesis of compounds of structure 22 containing an acylated amine functionality on the 6-position of the indole ring. These compounds can be synthesised from 6-nitro indoles of structure 18 in 4 steps. In the first step the indole of structure 18 is alkylated on the nitrogen atom by reaction of a halide of type R^1CH_2Y , in which Y is halogen, with NaH as a base in DMF to give compounds of structure 19. In the second step the nitro group of structure 19 is reduced to an amine group by $SnCl_2$ to give structure 20. Subsequent acylation of the amine group with an acid chloride of type $ZCOCl$ afforded structure 21, which was then sulfenylated at the 3-position of the indole ring by reaction with a sulfonyl chloride in either CH_2Cl_2 or Et_2O as a solvent, at room temperature, to give compounds of structure 22. Direct sulfenylation of a compound of structure 20 with a sulfonyl chloride afforded structure 23.

The present invention also relates to a pharmaceutical composition comprising the non-steroidal compound according to the invention mixed with a pharmaceutically acceptable auxiliary, such as described in the standard reference Gennaro *et al*, *Remington: The Science and Practice of Pharmacy*, (20th ed., Lippincott Williams & Wilkins, 2000, see especially Part 5: Pharmaceutical Manufacturing). Suitable auxiliaries are made available in e.g. the Handbook of Pharmaceutical Excipients (2nd Edition, Editors A. Wade and P.J. Weller; American Pharmaceutical Association;

Washington; The Pharmaceutical Press; London, 1994). The mixture of a compound according to the invention and a pharmaceutically acceptable auxiliary may be compressed into solid dosage units, such as pills, tablets, or be processed into capsules or suppositories. By means of pharmaceutically suitable liquids the compounds can also
5 be applied as an injection preparation in the form of a solution, suspension, emulsion, or as a spray, e.g. nasal spray. For making dosage units, e.g. tablets, the use of conventional additives such as fillers, colorants, polymeric binders and the like is contemplated. In general any pharmaceutically acceptable additive which does not interfere with the function of the active compounds can be used. The compounds of the
10 invention may also be included in an implant, a vaginal ring, a patch, a gel, and any other preparation for sustained release.

Suitable carriers with which the compositions can be administered include lactose, starch, cellulose derivatives and the like, or mixtures thereof used in suitable amounts.

15 The dosage amounts of the present compounds will be of the normal order for pharmaceutically active compounds, e.g. of the order of 0.001 to 50 mg/kg body weight of the recipient per administration. The recipient can be a human or an animal in need of an androgen receptor-related treatment.


20

The invention is illustrated by the following examples.

Examples

Example 1

- 5 1-(3,5-Difluoro-benzyl)-6-methoxy-3-(2-nitro-phenylsulfanyl)-1H-indole (Compound 63, Structure 3 of Scheme I, where $R^1 = 3,5$ -difluorophenyl, $R^2 = 2$ -nitrophenyl, $R^3 = R^4 = H$, $R^5 = OMe$)



10 **General method 1:** N-alkylation of a (un)substituted indole of structure 1 to give N-alkylated indole of structure 2, followed by 3-sulfenylation to give substituted indoles of structure 3 (Scheme I).

- (a) 1-(3,5-Difluoro-benzyl)-6-methoxy-1H-indole (Structure 2 of Scheme I, where $R^1 = 3,5$ -difluorophenyl, $R^3 = R^4 = H$, $R^5 = OMe$)

- 15 Under a nitrogen atmosphere: to a cooled (0°C) solution of 6-methoxyindole (863 mg, 5.86 mmol) in DMF (40 mL) was added NaH (60% in oil; 281 mg, 7.03 mmol) in small portions over a 3 minute period. The resulting green suspension was stirred at 0°C for 10 min. Then 3,5-difluorobenzyl bromide (0.91 mL, 7.03 mmol) was added. The mixture was stirred at 0°C for 1 h and then at room temperature for another 21 h. Ethyl acetate (50 mL) was added and the mixture was washed with 3% aqueous citric acid (3x50 mL) and brine (50 mL). The organic phase was dried (MgSO₄) and concentrated
20 *in vacuo* to give a green oil (1.43 g). The crude product was purified over a 20 g silica SPE cartridge (ethyl acetate/heptane 1:9) to give the title compound as a colourless oil (1.23 g, yield = 77%).

LCMS: 4.01 min (96.3%, $MH^+ = 274$); TLC (ethyl acetate/heptane 1:4): $R_f = 0.46$; 1H NMR (CDCl₃): δ 3.80 (s, 3H, OCH₃), 5.24 (s, 2H, NCH₂Ar), 6.51 (dd, 1H, $J_1 = 3.5$ Hz,

J2 = 0.8 Hz), 6.57-6.60 (m, 2H), 6.65 (d, 1H, J = 3.1 Hz), 6.66-6.72 (m, 1H), 6.81 (dd, 1H, J1 = 8.6 Hz, J2 = 3.1 Hz), 7.01 (d, 1H, J = 3.5 Hz), 7.53 (d, 1H, J = 8.6 Hz).

(b) 1-(3,5-Difluoro-benzyl)-6-methoxy-3-(2-nitro-phenylsulfanyl)-1H-indole

5 (Compound 63 . Structure 3 of Scheme I, where R¹ = 3,5-difluorophenyl, R² = 2-nitrophenyl, R³=R⁴=H, R⁵=OMe)

To a solution of 1-(3,5-Difluoro-benzyl)-6-methoxy-1H-indole (900 mg, 3.29 mmol) in diethyl ether (20 mL) was added dropwise at room temperature a suspension of 2-nitrobenzenesulfonyl chloride (627 mg, 3.31 mmol) in diethyl ether (10 mL) over a
10 period of 2 min. After stirring at room temperature for 1 h ethyl acetate (50 mL) was added and the mixture was washed with saturated NaHCO₃ solution (2x50 mL) and brine (50 mL). The organic phase was dried (MgSO₄) and concentrated *in vacuo* to give an orange-red oil (1.54 g). The crude product was crystallised from toluene/acetone to give the title compound as orange-red crystalline solid (900 mg, yield = 64%).

15 LCMS: 4.25 min (100%, MH⁺ = 427); HPLC: 4.86 min (98.7%); ¹H NMR (CDCl₃): δ 3.82 (s, 3H, OCH₃), 5.32 (s, 2H, NCH₂Ar), 6.63-6.69 (m, 2H), 6.72-6.79 (m, 1H), 6.75 (d, 1H, J = 2.7 Hz), 6.85 (dd, 1H, J1 = 8.2 Hz, J2 = 2.7 Hz), 6.98 (dd, 1H, J1 = 6.7 Hz, J2 = 1.2 Hz), 7.16-7.20 (m, 1H), 7.26-7.30 (m, 1H) 7.34 (s, 1H), 7.39 (d, 1H, J = 8.2 Hz), 8.27 (dd, 1H, J1 = 8.2 Hz, J2 = 1.6 Hz).

20 According to General method 1 the following compounds were prepared:




Table 1 Compounds synthesised according to General Method 1


Compound X							LCMS	LCMS
								ref.
number		R ¹	R ²	R ³	R ⁴	R ⁵	MWt (MH ⁺)	Time ^a (min)
21	S	3,5-difluorophenyl	2-methoxyphenyl	H	H	OMe	411.47	446 ^a 5.14 ^b
22	S	3,5-difluorophenyl	2-nitrophenyl	H	H	Br	475.32	476 4.89
23	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CF ₃	464.41	465 5.34 ^b
24	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CH ₂ OH	426.44	Nd 3.88 ^b

Compound X							LCMS	LCMS
							(MH ⁺)	ret. Time ^a
number	R ¹	R ²	R ³	R ⁴	R ⁵	MWt	(MH ⁺)	(min)
25	S 3,5-difluorophenyl	2-nitrophenyl	H	H	Cl	430.86	431	4.89
50	S 3,5-difluorophenyl	2-nitrophenyl	H	H	F	414.40	415	4.79
51	S 3,5-difluorophenyl	2-nitrophenyl	F	H	H	414.40	415	4.72
52	S 3,5-difluorophenyl	2-nitrophenyl	H	H	H	396.41	397	4.22
53	S 3,5-difluorophenyl	2-nitrophenyl	H	F	H	414.40	415	4.73
54	S 3,5-difluorophenyl	2-nitrophenyl	Cl	H	H	430.86	431	5.14 ^b
55	S 3,5-difluorophenyl	2-nitrophenyl	Me	H	H	410.44	411	5.24 ^b
56	S 3,5-difluorophenyl	2-nitrophenyl	H	OH	H	412.41	413	4.50 ^b
60	S 3,5-difluorophenyl	2-nitrophenyl	H	H	NO ₂	441.41	442	4.7
61	S 3,5-difluorophenyl	2-nitrophenyl	H	H	OH	412.41	413	4.54 ^b
63	S 3,5-difluorophenyl	2-nitrophenyl	H	H	OMe	426.45	274 ^a	4.01
66	S 3,5-difluorophenyl	2-pyridyl-N-oxide	H	H	OMe	398.43	399	3.96
90	S phenyl	2-nitrophenyl	H	H	H	360.43	361	4.19
91	S phenyl	2-nitrophenyl	H	OMe	H	390.46	391	4.14
92	S phenyl	2-nitrophenyl	H	H	OMe	390.46	391	4.12

a) 7 min LCMS method was used, unless stated otherwise. b) 10 min LCMS method was used. c) M+Cl. nd = not detected. d) = $[M-S(PhNO_2)]H^+$.

Example 2

- 5 1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1*H*-indole-6-carboxylic acid methyl ester (Compound 31, Structure 3 of Scheme I, where R¹ = 3,5-difluorophenyl, R² = 2-nitrophenyl, R³=R⁴=H, R⁵=CO₂Me)



(a) 1*H*-Indole-6-carboxylic acid methyl ester (Structure 1 of Scheme I, where $R^3=R^4=H$, $R^5=CO_2Me$)

To a solution of 1*H*-Indole-6-carboxylic acid (1.500 g, 9.31 mmol) in methanol (50 mL) was added concentrated H_2SO_4 (550 μ L, 10.24 mmol). The mixture was stirred overnight at reflux temperature. The mixture was then neutralised to pH 7 by addition of saturated aqueous $NaHCO_3$ and the mixture was extracted twice with ethyl acetate. The combined organic layers were dried over Na_2SO_4 and concentrated *in vacuo* to give a yellow powder. The product was recrystallised from heptane/ethyl acetate to give the title compound as green/yellow crystals (688 mg, yield = 53%)

10. 1H NMR ($CDCl_3$): δ 3.94 (s, 3H, CH_3OCO), 6.60 (m, 1H), 7.37 (t, 1H, $J = 4.7$ Hz), 7.66 (d, 1H, $J = 8.2$ Hz), 7.82 (dd, 1H, $J_1 = 8.2$ Hz, $J_2 = 1.0$ Hz), 8.18 (s, 1H), 8.73 (s, 1H, NH)

(b) 1-(3,5-Difluoro-benzyl)-1*H*-indole-6-carboxylic acid methyl ester (Structure 2 of Scheme I, where $R^1 = 3,5$ -difluorophenyl, $R^3=R^4=H$, $R^5=CO_2Me$)

Under a nitrogen atmosphere: to a solution of 1*H*-indole-6-carboxylic acid methyl ester (367 mg, 2.09 mmol) in DMF (10 mL) was added NaH (60% in oil, 101 mg, 2.52 mmol) at room temperature. After stirring for 15 min 1-bromomethyl-3,5-difluorobenzene (325 μ L, 2.51 mmol) was added and the mixture was stirred at room temperature for 4 h. The reaction was quenched with 3% aqueous citric acid (10 mL) and ethyl acetate (30 mL) was added. The mixture was washed with 3% aqueous citric acid (3x20 mL) and brine (20 mL). The organic phase was dried ($MgSO_4$) and concentrated *in vacuo* to give a pale yellow oil (702 mg). The crude product was purified over a 20 g silica SPE cartridge (ethyl acetate/heptane 1:9) to give the title compound as a colourless oil, which slowly crystallised on standing (530 mg, yield = 84%).

LCMS: 4.08 min (99%, $MH^+ = 302$); 1H NMR ($CDCl_3$): δ 3.92 (s, 3H, CO_2CH_3), 5.37 (s, 2H, NCH_2Ar), 6.54-6.60 (m, 2H), 6.64 (d, 1H, $J = 3.1$ Hz), 6.67-6.74 (m, 1H), 7.27 (d, 1H, $J = 3.1$ Hz), 7.68 (d, 1H, $J = 8.6$ Hz), 7.83 (d, 1H, $J = 8.6$ Hz), 8.01 (s, 1H).


(c) 1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1*H*-indole-6-carboxylic acid methyl ester (Compound 31, Structure 3 of Scheme I, where $R^1 = 3,5$ -difluorophenyl, $R^2 = 2$ -nitrophenyl, $R^3=R^4=H$, $R^5=CO_2Me$)

To a solution of 1-(3,5-Difluoro-benzyl)-1*H*-indole-6-carboxylic acid methyl ester (330 mg, 1.10 mmol) in dichloromethane (30 mL) was added at room temperature a solution of 2-nitrobenzenesulfonyl chloride (210 mg, 1.11 mmol) in dichloromethane (10 mL). The mixture was stirred at room temperature for 4 days. The reaction mixture was concentrated and the crude product was purified over a 20 g silica SPE cartridge (ethyl acetate/heptane 1:5 to 1:2) to give the title compound as a yellow solid (392 mg, yield = 78%).

HPLC: 4.60 min (97.8%); ¹H NMR (CDCl₃): δ 3.93 (s, 3H, CO₂CH₃), 5.45 (s, 2H, NCH₂Ar), 6.63-6.69 (m, 2H), 6.74-6.80 (m, 1H), 6.88 (dd, 1H, J₁ = 8.2 Hz, J₂ = 1.2 Hz), 7.18-7.23 (m, 1H), 7.26-7.30 (m, 1H), 7.57 (d, 1H, J = 7.8 Hz), 7.58 (s, 1H), 7.89 (dd, 1H, J₁ = 7.8 Hz, J₂ = 1.2 Hz), 8.12 (s, 1H), 8.28 (dd, 1H, J₁ = 7.8 Hz, J₂ = 1.2 Hz).

Example 3

1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1*H*-indole-6-carboxylic acid
(Compound 30, Structure 14 of Scheme IV, where R¹ = 3,5-difluorophenyl, R² = 2-nitrophenyl, R³=R⁴=H)



To a solution of 1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1*H*-indole-6-carboxylic acid methyl ester (150.0 mg, 0.33 mmol) in dioxane (20 mL) and water (15 mL) was added LiOH.H₂O (83.1 mg, 2.0 mmol) in 5 ml water. The reaction mixture was stirred overnight at 60°C. The mixture was then acidified to pH 4 by addition of 15% aqueous HCl and the mixture was extracted twice with ethyl acetate. The combined organic layers were dried over Na₂SO₄ and concentrated *in vacuo* to give a yellow powder. The product was recrystallised from heptane/ethyl acetate to give the title compound as yellow/orange crystals (129.3 mg, y = 89%).


¹H NMR (DMSO): δ 5.62 (s, 2H, CH₂Ar), 6.85 (dd, 1H, J₁ = 8.6 Hz, J₂ = br), 7.05 (m, 2H, br), 7.20 (m, 1H, br), 7.32-7.38 (m, 1H), 7.40 (d, 1H, J = 8.2 Hz), 7.44-7.50 (m, 1H), 7.68 (dd, 1H, J₁ = 8.6 Hz, J₂ = br), 7.97 (s, 1H, COOH), 8.20 (s, 1H), 8.26 (s, 1H), 8.28 (d, 1H, J = 8.2)

5

Example 4

1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1H-indole-6-carboxylic acid methylamide (Compound 47, Structure 15 of Scheme IV, where R¹ = 3,5-difluorophenyl, R² = 2-nitrophenyl, R³=R⁴=R⁸=H, R⁹=Me)

10



General method 2: amidation of 6-carboxyl indoles of structure 14 to give 6-carboxamideindoles of structure 15 (Scheme IV).

Under nitrogen atmosphere: to a solution of 1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1H-indole-6-carboxylic acid (25.1 mg, 57.0 μmol) in dry DMF (10mL) was added NMM (1 mL), HOBT (9.60 mg, 62.7 μmol), EDCI (12.1 mg, 62.7 μmol), and methylammonium chloride (19.2 mg, 285 μmol). The reaction mixture was stirred overnight at room temperature and then poured into ice water. The resulting precipitate was filtered and the residue was washed with excess water followed by little heptane. The product was dried *in vacuo* at 40°C to give the title compound as a yellow solid (15.2 mg, y = 60%).

LCMS: 4.31 min (100.0%, MH⁺ = 454); ¹H NMR (DMSO): δ 2.80(d, 3H, J = 4.3, CH₃NHCO), 5.63 (s, 2H, CH₂Ar), 6.86 (dd, 1H, J₁ = 8.2 Hz, J₂ = 1.0 Hz), 7.04 (m, 2H), 7.20 (m, 1H), 7.33-7.37 (m, 1H), 7.41 (d, 1H, 8.6 Hz), 7.46-7.50 (m, 1H), 7.64 (dd, 1H, J₁ = 8.2 Hz, J₂ = 1.0 Hz), 8.15 (s, 1H), 8.25 (s, 1H), 8.27 (dd, 1H, J₁ = 8.6 Hz, J₂ = 1.0 Hz), 8.40 (m, 1H, MeNHCO).

According to General method 2 the following compounds were prepared:




Table 2 Compounds synthesised according to General Method 2


Compound X		LCMS		LCMS		LCMS		ret. Time ^a (min)
number	R ¹	R ²	R ³	R ⁴	R ⁵	MWt	(MH ⁺)	
27	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CO(1-pyrrolidinyl)	493.53	494	4.66
28	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CO(4-morpholinyl)	509.53	510	4.50
29	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CO[1-(4-methylpiperazinyl)]	522.57	523	3.81
32	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONEt ₂	495.55	496	4.82
34	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ (2-furanyl)	519.53	520	4.74
35	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ (3-pyridyl)	530.55	531	3.94
36	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ CH ₂ NMe ₂	510.56	511	3.90
37	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ CH ₂ OH	483.49	484	4.21
38	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ CH ₂ OMe	497.52	498	4.48
39	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ CO ₂ Me	511.50	512	4.51
40	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ CONMe ₂	524.55	525	4.33
41	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ cPr	493.53	494	4.75
42	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ iPr	495.55	496	4.85
43	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ Ph	529.56	530	4.90
44	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHcPr	479.50	480	4.57
45	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHEt	467.49	468	4.56
46	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHiPr	481.52	482	4.71
47	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHMe	453.47	454	4.31
48	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONHnPr	481.52	482	4.75
49	S 3,5-difluorophenyl	2-nitrophenyl	H	H	CONMe ₂	467.49	468	4.30 ^b

a) 7 min LCMS method was used, unless stated otherwise. b) 10 min LCMS method

5 was used.

Example 5

1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1*H*-indole-6-carboxylic acid amide
 (Compound 33, Structure 16 of Scheme IV, where $R^1 = 3,5$ -difluorophenyl, $R^2 = 2$ -nitrophenyl, $R^3 = R^4 = H$)



5

The title compound was prepared according to General method 2, using 51.1 mg (116 μ mol) carboxylic acid, 19.5 mg (128 μ mol) HOBt, 24.6 mg (128 μ mol) EDCI and 31.1 mg (581 μ mol) ammonium chloride. The title compound was obtained as a yellow solid (9.0 mg; 18%).


LCMS: 4.16 min (100.0%, $MH^+ = 440$); 1H NMR ($CDCl_3$): δ 5.46 (s, 2H, CH_2Ar), 6.65 (m, 2H), 6.77 (m, 1H), 6.88 (dd, 1H, $J_1 = 8.2$ Hz, $J_2 = 1.0$ Hz), 7.21 (m, 1H), 7.29 (m, 1H), 7.51 (dd, 1H, $J_1 = 7.8$ Hz, $J_2 = 1.0$ Hz), 7.57 (d, 1H, $J = 7.7$ Hz), 7.58 (s, 1H), 8.05 (s, 1H), 8.29 (dd, 1H, $J_1 = 8.2$ Hz, $J_2 = 1.0$ Hz).

15

Example 6

6-Methoxy-3-(2-nitro-phenylsulfanyl)-1-pyridin-2-ylmethyl-1*H*-indole (Compound 9, Structure 3 of Scheme I, where $R^1 = 2$ -pyridyl, $R^2 = 2$ -nitrophenyl, $R^3 = R^4 = H$, $R^5 = OMe$)

20



General method 3: 3-sulfenylation of a (un)substituted indole of structure 1 to give substituted indoles of structure 4, followed by indole-*N*-alkylation to give *N*-alkylated indole structure 3 (Scheme I).

- 5 (a) 6-Methoxy-3-(2-nitro-phenylsulfanyl)-1*H*-indole (Structure 4 of Scheme I, where $R^2 = 2\text{-nitrophenyl}$, $R^1=R^3=R^4=H$, $R^5=OMe$)

To a solution of 6-methoxyindole (1.5 g, 10.2 mmol) in Et_2O (100 mL) was added a solution of 2-nitrobenzenesulfonylchloride (1.93 g, 10.2 mmol) in 50 mL Et_2O dropwise, over a 4 minute period. The resulting yellow solution was stirred at room temperature for 1 h. The solvent was evaporated and the crude product was purified
10 over a silica column (heptane/ethyl acetate 9:1) to give the title compound (3.054 g, yield = 97%).

HPLC : 3.72 min, purity 96.7%, TLC (heptane/ethyl acetate 1:1): $R_f = 0.6$;

1H NMR ($CDCl_3$): δ 3.87 (s, 3H, OCH_3), 6.84 (dd, 1H, $J_1 = 7.8$ Hz, $J_2 = 3.13$ Hz), 6.97 (m, 2H), 7.16 (t, 1H, $J = 7.8$ Hz), 7.25 (t, 1H, $J = 7.8$ Hz), 7.36 (d, 1H, $J = 7.8$ Hz), 7.44
15 (d, 1H, $J = 2.0$ Hz), 8.26 (d, 1H, $J = 7.8$ Hz), 8.45 (s, 1H, NH).

- (b) 6-Methoxy-3-(2-nitro-phenylsulfanyl)-1-pyridin-2-ylmethyl-1*H*-indole (Compound 9, Structure 3 of Scheme I, where $R^1 = 2\text{-pyridyl}$, $R^2 = 2\text{-nitrophenyl}$, $R^3=R^4=H$, $R^5=OMe$)


20 To a solution of 6-Methoxy-3-(2-nitro-phenylsulfanyl)-1*H*-indole (1.5 g, 5 mmol) in DMF (150 mL) was added NaH (60% in oil; 500 mg, 12.5 mmol) in small portions, over a 4 minute period. The resulting dark solution was stirred for 5 min. at room temperature. Then 2-picolyl chloride hydrochloride (984 mg, 6 mmol) was added in small portions, over a 2 minute period. During stirring at room temperature (2h) the
25 colour of the solution slightly changed from dark to yellow. Ethyl acetate (100 mL) was added and the mixture was washed with 2% aqueous citric acid (2x100 mL) and water (100 mL). The organic phase was dried ($MgSO_4$) and the solvent was evaporated. The crude product was purified by crystallisation (ethyl acetate/heptane) to give the title compound (1.437 g, yield = 74%).

30 HPLC : 10.3 min, purity 99.6%, TLC (heptane/ethyl acetate 1:1): $R_f = 0.3$;

1H NMR ($CDCl_3$): δ 3.80 (s, 3H, OCH_3), 5.47 (s, 2H, NCH_2), 6.82 (m, 2H), 6.88 (d, 1H, $J = 7.8$ Hz), 7.01 (dd, 1H, $J_1 = 7.8$ Hz, $J_2 = 1.6$ Hz), 7.15-7.19 (m, 1H), 7.25 (m,

2H), 7.37 (d, 1H, $j = 8.6$ Hz), 7.60-7.64 (m, 1H), 8.26 (dd, 1H, $J_1 = 7.8$ Hz, $J_2 = 1.6$ Hz), 8.63 (d, 1H, $J = 5.88$ Hz).

According to General method 3 the following compounds were prepared:



5 Table 3 Compounds synthesised according to General Method 3


Compound								LCMS	LCMS ret.
number	X	R ¹	R ²	R ³	R ⁴	R ⁵	MWt	(MH ⁺)	Time ^a (min)
1	S	2,5-difluorophenyl	2-nitrophenyl	H	H	OMe	426.44	427	4.07
2	S	2,fluoro,3-methylphenyl	2-nitrophenyl	H	H	OMe	422.48	423	4.23
3	S	2-chlorophenyl	2-nitrophenyl	H	H	OMe	424.91	425	4.17
4	S	2-cyanophenyl	2-nitrophenyl	H	H	OMe	415.47	416	3.97
5	S	2-fluorophenyl	2-nitrophenyl	H	H	OMe	408.45	409	4.10
6	S	2-methyl,3-nitrophenyl	2-nitrophenyl	H	H	OMe	449.48	450	4.10
7	S	2-methylphenyl	2-nitrophenyl	H	H	OMe	404.49	405	4.16
8	S	2-nitrophenyl	2-nitrophenyl	H	H	OMe	435.46	436	4.08
9	S	2-pyridyl	2-nitrophenyl	H	H	OMe	391.45		10.30 ^b
11	S	2-tetrahydropyranyl	2-nitrophenyl	H	H	OMe	398.48	399	4.68
12	S	2-trifluoromethylphenyl	2-nitrophenyl	H	H	OMe	458.46	459	4.29
13	S	3-(5-methylisoxazolyl)	2-nitrophenyl	H	H	OMe	395.44	396	4.66
14	S	3,4-dichlorophenyl	2-nitrophenyl	H	H	OMe	459.35	459	4.33
15	S	3,5-dichlorophenyl	2-nitrophenyl	H	H	OMe	459.35	459	4.36
67	S	3-chlorophenyl	2-nitrophenyl	H	H	OMe	424.91	425	4.34
68	S	3-cyanophenyl	2-nitrophenyl	H	H	OMe	415.47	416	3.92
69	S	3-fluoro,5-trifluoromethylphenyl	2-nitrophenyl	H	H	OMe	476.45	477	4.42
70	S	3-fluoro,6-trifluoromethylphenyl	2-nitrophenyl	H	H	OMe	476.45	477	4.27
71	S	3-fluorophenyl	2-nitrophenyl	H	H	OMe	408.45	409	4.22
72	S	3-methoxyphenyl	2-nitrophenyl	H	H	OMe	420.49	421	4.04
73	S	3-methylphenyl	2-nitrophenyl	H	H	OMe	404.49	405	4.16
74	S	3-nitrophenyl	2-nitrophenyl	H	H	OMe	435.46	436	4.00

Compound							LCMS		LCMS ret.
number	X	R ¹	R ²	R ³	R ⁴	R ⁵	MWt	(MH ⁺)	Time ^a (min)
75	S	3-pyridyl	2-nitrophenyl	H	H	OMe	391.45	392	3.28
76	S	3-trifluoromethyl,4-chlorophenyl	2-nitrophenyl	H	H	OMe	492.90	493	4.33
77	S	3-trifluoromethoxyphenyl	2-nitrophenyl	H	H	OMe	474.46	475	4.43
78	S	3-trifluoromethylphenyl	2-nitrophenyl	H	H	OMe	458.46	459	4.21
79	S	4-(2-methylthiazolyl)	2-nitrophenyl	H	H	OMe	411.50	412	4.39
80	S	4-(3,5-dimethylisoxazolyl)	2-nitrophenyl	H	H	OMe	409.46	410	4.49
81	S	4-chlorophenyl	2-nitrophenyl	H	H	OMe	424.91	425	4.22
82	S	4-cyanophenyl	2-nitrophenyl	H	H	OMe	415.47	416	3.93
83	S	4-fluorophenyl	2-nitrophenyl	H	H	OMe	408.45	409	4.07
84	S	4-methoxyphenyl	2-nitrophenyl	H	H	OMe	420.49	421	4.03
85	S	4-methylphenyl	2-nitrophenyl	H	H	OMe	404.49	405	4.19
86	S	4-morpholinyl	2-nitrophenyl	H	H	OMe	413.50	414	3.54
87	S	5-(2-chlorothiazolyl)	2-nitrophenyl	H	H	OMe	431.92	432	4.61
88	S	5-(2-chlorothiophenyl)	2-nitrophenyl	H	H	OMe	430.93	431	4.92
89	S	cyclohexyl	2-nitrophenyl	H	H	OMe	396.51	397	5.11

a) 7 min LCMS method was used unless stated otherwise. b) a 20 min run on an HPLC system was used.

Example 7

- 5 1-(3,5-Difluoro-benzyl)-6-methoxy-3-(pyridin-2-ylsulfanyl)-1H-indole (Compound 65, Structure 3 of Scheme I, where R¹ = 3,5-difluorophenyl, R² = 2-pyridyl, R³=R⁴=H, R⁵=OMe)




Under a nitrogen atmosphere: to a cooled (-10°C) solution of 2-mercaptopyridine (25 mg, 0.22 mmol) in CCl₄ (2 mL) was passed Cl₂-gas during a period of 1 minute. The solvent was evaporated and a solution of 1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indole (25 mg, 0.09 mmol) in Et₂O (2 mL) was added dropwise. The reaction mixture was concentrated and the crude product was purified by preparative LCMS to give the title compound (7.77 mg, yield = 23%).

LCMS : 3.95 min (100%, MH⁺ 383); TLC (heptane/ethyl acetate 1:1): R_f = 0.6;

¹H-NMR (CDCl₃) : δ 2.23 (s, 1H, NH), 3.82 (s, 1H, OCH₃), 6.65 (m, 2H), 6.73 (d, 1H, J = 2.7 Hz), 6.73-6.78 (m, 1H), 6.80 (d, 1H, J = 7.8 Hz), 6.87 (dd, 1H, J₁ = 7.8 Hz, J₂ = 1.6 Hz), 7.00 (ddd, 1H, J₁ = 9.8 Hz, J₂ = 4.7 Hz, J₃ = 0.8 Hz), 7.35 (s, 1H), 7.42 (ddd, 1H, J₁ = J₂ = 7.8 Hz, J₃ = 1.6 Hz), 7.50 (d, 1H, 8.6 Hz), 8.46 (d, 1H, J = 4.7 Hz).

Example 8

2-[1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indol-3-ylsulfanyl]-phenyl-methanol
(Compound 16, Structure 8 of Scheme III, where R¹ = 3,5-difluorophenyl, R³=R⁴=H, R⁵=OMe)



a) 2-[1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indol-3-ylsulfanyl]-benzoic acid methyl ester
(Compound 19, Structure 7 of Scheme III, where R¹ = 3,5-difluorophenyl, R³=R⁴=H, R⁵=OMe)

Chlorine gas was bubbled through CCl₄ (5 mL) at -10°C for 3 min. Then a solution of methylthiosalicylate (91 µL, 0.66 mmol) in CCl₄ (2 mL) was added. The mixture was stirred at -10°C for 5 min and then at room temperature for 15 min. The mixture was concentrated and redissolved in CH₂Cl₂ (5 mL). To this solution a solution of 1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indole (121 mg, 0.443 mmol) in CH₂Cl₂ (3 mL) was

added dropwise. The mixture was stirred at room temperature for 2 h and was concentrated. The crude product was purified by column chromatography (ethyl acetate/heptane 1:9) to give the title compound as a white solid (161 mg, 83% yield).

LCMS: 4.72 min ($MH^+ = 440$); 1H -NMR ($CDCl_3$): δ 3.82 (s, 3H, $ArOCH_3$), 3.99 (s, 3H, CO_2CH_3), 5.30 (s, 2H, CH_2Ar), 6.62-6.69 (m, 2H), 6.71-6.77 (m, 2H), 6.83 (dd, 1H, $J_1 = 8.6$ Hz, $J_2 = 2.4$ Hz), 6.87 (dd, 1H, $J_1 = 8.4$ Hz, $J_2 = 1.2$ Hz), 7.06-7.11 (m, 1H, 7.15-7.20 (m, 1H), 7.30 (s, 1H), 7.44 (d, 1H, $J = 8.6$ Hz), 8.02 (dd, 1H, $J_1 = 8.0$ Hz, $J_2 = 1.2$ Hz).


- 10 b) 2-[1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indol-3-ylsulfanyl]-phenyl}-methanol (Compound 16, Structure 8 of Scheme III, where $R^1 = 3,5$ -difluorophenyl, $R^3=R^4=H$, $R^5=OMe$)

Under a nitrogen atmosphere: to a solution of 2-[1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indol-3-ylsulfanyl]-benzoic acid methyl ester (13.5 mg, 0.031 mmol) in THF (2 mL) was added $LiAlH_4$ (1.0 M in THF, 0.040 mL, 0.040 mmol). The mixture was stirred at room temperature for 3 h. Then ethyl acetate (25 mL) was added and the mixture was washed with 3% aqueous citric acid (2x25 mL) and brine (25 mL). The organic phase was dried ($MgSO_4$) and concentrated to give a colourless oil (15 mg). The crude product was purified by prep. HPLC to give the title compound as a colourless oil (5.6 mg, 44% yield).

LCMS: 4.47 min (95.6%, $MH^+ = 412$); 1H -NMR ($CDCl_3$): δ 3.49 (s, 1H, OH), 3.81 (3, 3H, OCH_3), 4.92 (s, 2H, CH_2OH), 5.28 (s, 2H, CH_2Ar), 6.62-6.66 (m, 2H), 6.70-6.77 (m, 2H), 6.83 (dd, 1H, $J_1 = 8.8$ Hz, $J_2 = 2.4$ Hz), 6.89 (dd, 1H, $J_1 = 8.0$ Hz, $J_2 = 1.2$ Hz), 7.02-7.07 (m, 1H), 7.08-7.13 (m, 1H), 7.29 (s, 1H), 7.38 (dd, 1H, $J_1 = 8.0$ Hz, $J_2 = 1.2$ Hz), 7.44 (d, 1H, $J = 8.8$ Hz).

Example 9

2-[1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indol-3-ylsulfanyl]-benzamide (Compound 18, Structure 11 of Scheme III, where $R^1 = 3,5$ -difluorophenyl, $R^3=R^4=H$, $R^5=OMe$)



5

a) 2-[1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indol-3-ylsulfanyl]-benzoic acid (Structure 9 of Scheme III, where $R^1 = 3,5$ -difluorophenyl, $R^3=R^4=H$, $R^5=OMe$)

To a solution of 2-[1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indol-3-ylsulfanyl]-benzoic acid methyl ester (97 mg, 0.221 mmol) in THF (10 mL) was added a solution of

10 LiOH·H₂O (71 mg, 1.69 mmol) in water (10 mL). The mixture was stirred at room temperature for 17 h and then at 60°C for 24 h and was then acidified with 3% aqueous citric acid. Ethyl acetate (50 mL) was added and the mixture was washed with 3% aqueous citric acid (2x50 mL) and brine (50 mL). The organic phase was dried (MgSO₄) and concentrated to give a yellow solid (104 mg). The crude product was

15 recrystallised from ethyl acetate/heptane to give the title compound as a yellow powder (49 mg, 52% yield).

LCMS: 4.37 min ($MH^+ = 426$); ¹H-NMR (CDCl₃): δ 3.77 (s, 3H, OCH₃), 5.51 (s, 2H, CH₂Ar), 6.69 (d, 1H, J = 8.0 Hz), 6.76 (dd, 1H, J₁ = 8.4 Hz, J₂ = 2.4 Hz), 6.98-7.04 (m, 2H), 7.11-7.15 (m, 1H), 7.16-7.25 (m, 4H), 7.83 (s, 1H), 7.92 (dd, 1H, J₁ = 8.0 Hz, J₂ = 1.6 Hz).

20

General method 4: Reaction of an amine with a carboxylic acid of structure 9, with TBTU and DIPEA, to give amides of structure 3, in which $R^2 =$ phenyl-2-carboxamide, exemplified by compounds of structure 10 and 11 (Scheme III).

25


b) 2-[1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indol-3-ylsulfanyl]-benzamide
(Compound 18, Structure 11 of Scheme III, where $R^1 = 3,5$ -difluorophenyl, $R^3=R^4=H$, $R^5=OMe$)

Through a solution of 2-[1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indol-3-ylsulfanyl]-
benzoic acid (36 mg, 0.085 mmol), TBTU (28 mg, 0.087 mmol) and DIPEA (30 μ L,
0.172 mmol) in DMF (1 mL) was bubbled NH_3 gas for 10 min. The mixture was then
stirred at room temperature for 3 days. Then 3% aqueous citric acid (0.4 mL) was added
and the crude reaction mixture was purified over a RP-SPE cartridge (2g sorbent; 25%
aqueous methanol to 100% methanol) to give a white solid (34 mg). The crude product
was purified by column chromatography (ethyl acetate) to give the title compound as a
colourless oil, which solidifies on standing (31 mg, yield: 86%).

LCMS: 4.20 min ($MH^+ = 425$); 1H -NMR ($CDCl_3$) : δ 3.81 (s, 3H, OCH_3), 5.39 (s, 2H, CH_2Ar), 6.05 (s, br, 1H, NH), 6.28 (s, br, 1H, NH), 6.61-6.67 (m, 2H), 6.70-6.72 (m, 1H), 6.73-6.77 (m, 1H), 6.83 (dd, 1H, $J_1 = 8.8$ Hz, $J_2 = 2.0$ Hz), 6.92 (dd, 1H, $J_1 = 8.0$ Hz, $J_2 = 1.2$ Hz), 7.07-7.17 (m, 2H), 7.31 (s, 1H), 7.45 (d, 1H, $J = 8.8$ Hz), 7.60 (dd, 1H, $J_1 = 8.0$ Hz, $J_2 = 1.2$ Hz).

Example 10


2-[1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indol-3-ylsulfanyl]-*N*-methyl-benzamide
(Compound 17, Structure 10 of Scheme III, where $R^1 = 3,5$ -difluorophenyl, $R^3=R^4=H$, $R^5=OMe$)



According to general method 4 compound 17 was synthesised from methylamine.
LCMS: 4.07 min ($MH^+ = 439$),

Example 11


2-[1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indol-3-ylsulfanyl]-benzonitrile (Compound 20, Structure 12 of Scheme III, where $R^1 = 3,5$ -difluorophenyl, $R^3=R^4=H$, $R^5=OMe$)



- To a cooled (0°C) solution of 2-[1-(3,5-Difluoro-benzyl)-6-methoxy-1*H*-indol-3-ylsulfanyl]-benzamide (8 mg, 0.019 mmol) and Et₃N (10 µL, 0.072 mmol) in CH₂Cl₂ (1.5 mL) was added TF₂O (6 µL, 0.036 mmol). The mixture was stirred at 0°C for 2 h and then at room temperature for 22 h. Water (25 mL) was added and the mixture was extracted with CH₂Cl₂ (2x25 mL). The combined organic phases were dried (Na₂SO₄) and concentrated to give a brown oil (25 mg). The crude product was purified by column chromatography (ethyl acetate/heptane 1:2) to give the title compound as a pale pink oil (3 mg, yield: 39%).
- LCMS: 4.57 min (MH⁺ = 407); ¹H-NMR (CDCl₃): δ 3.82 (s, 3H, OCH₃), 5.30 (s, 2H, CH₂Ar), 6.62-6.67 (m, 2H), 6.71-6.73 (m, 1H), 6.74-6.78 (m, 1H), 6.86 (dd, 1H, J₁ = 8.8 Hz, J₂ = 2.0 Hz), 6.91 (d, 1H, J = 8.4 Hz), 7.11-7.15 (m, 1H), 7.26-7.30 (m, 1H), 7.37 (s, 1H), 7.46 (d, 1H, J = 8.8 Hz), 7.57-7.60 (m, 1H).

Example 12

1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1*H*-indole-6-carbonitrile
 (Compound 26, Structure 17 of Scheme IV, where R¹ = 3,5-difluorophenyl, R² = 2-nitrophenyl, R³=R⁴=H)




To a cooled (0°C) suspension of 1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1*H*-indole-6-carboxylic acid amide (15 mg, 0.034 mmol) and Et₃N (10 µL, 0.072 mmol) in CH₂Cl₂ (1.5 mL) was added Tf₂O (13 µL, 0.077 mmol). The mixture was stirred at 0°C for 2 h and then at room temperature for 20 h. Water (25 mL) was added and the
 5 mixture was extracted with CH₂Cl₂ (2x25 mL). The combined organic phases were dried (Na₂SO₄) and concentrated to give the title compound as a yellow solid (13 mg, yield: 91%).

LCMS: 4.55 min (MH⁺ = not detectable); ¹H-NMR (CDCl₃): δ 5.42 (s, 2H, CH₂Ar), 6.63-6.69 (m, 2H), 6.78-6.85 (m, 2H), 7.21-7.25 (m, 1H), 7.28-7.33 (m, 1H), 7.44 (dd, 1H, J₁ = 8.4 Hz, J₂ = 1.2 Hz), 7.62 (d, 1H, J = 8.4 Hz), 7.66 (s, 1H), 8.29 (dd, 1H, J₁ = 8.4 Hz, J₂ = 1.6 Hz).

Example 13

1-(3,5-Difluoro-benzyl)-6-methoxy-3-(2-nitro-benzenesulfinyl)-1*H*-indole (Compound
 15 64, Structure 5 of Scheme II, where R¹ = 3,5-difluorophenyl, R² = 2-nitrophenyl, R³=R⁴=H, R⁵=OMe)




To a solution of 1-(3,5-Difluoro-benzyl)-6-methoxy-3-(2-nitro-phenylsulfanyl)-1*H*-indole (40 mg, 0.09 mmol) in CH₂Cl₂ (4 mL) was added meta-chloroperbenzoic acid
 20 (14.5, 0.08 mmol). The solution was stirred at room temperature for 2 h. The solvent was evaporated and the crude product was purified by prep LC-MS to give the title compound (14.3 mg, yield = 34%).

HPLC: 4.13 min, purity 100%; TLC (heptane/ethyl acetate 1:1): R_f = 0.4; ¹H NMR (CDCl₃): δ 3.66 (s, 3H, OCH₃), 5.12 (s, 2H), 6.57 (m, 3H), 6.75 (m, 2H), 7.47 (d, 1H, J = 7.8 Hz), 7.52 (s, 1H), 7.66-7.70 (m, 1H), 8.28-8.32 (m, 1H), 8.23 (dd, 1H, J₁ = 7.8 Hz, J₂ = 1.9 Hz), 8.80 (dd, 1H, J₁ = 7.8 Hz, J₂ = 3.1 Hz).

Example 14

1-(3,5-Difluoro-benzyl)-6-methoxy-3-(2-nitro-benzenesulfonyl)-1*H*-indole (Compound 62, Structure 6 of Scheme II, where $R^1 = 3,5$ -difluorophenyl, $R^2 = 2$ -nitrophenyl, $R^3 = R^4 = H$, $R^5 = OMe$)



5

To a solution of 1-(3,5-Difluoro-benzyl)-6-methoxy-3-(2-nitro-phenylsulfanyl)-1*H*-indole (40 mg, 0.09 mmol) in CH_2Cl_2 (4 mL) was added meta-chloroperbenzoic acid (46.5 mg, 0.27 mmol). The solution was stirred at room temperature for 6 h. The solvent was evaporated and the crude product was purified by LC-MS to give the title compound (14.5 mg, yield = 34%).


10

HPLC: 4.38 min. purity 100%; TLC (heptane/ethyl acetate 1:1) : $R_f = 0.6$; 1H NMR ($CDCl_3$): δ 3.71 (s, 3H, OCH_3), 5.25 (s, 2H), 6.58 (d, 1H, $J = 3.1$), 6.62 (m, 1H), 6.07-6.11 (m, 1H), 6.88 (dd, 1H, $J_1 = 7.8$ Hz, $J_2 = 3.1$ Hz), 7.63 (m, 4H), 7.72 (d, 1H, $J = 7.8$ Hz), 7.83 (s, 1H), 8.37 (dd, 1H, $J_1 = 7.8$ Hz, $J_2 = 1.9$ Hz).

15

Example 15

2-[6-Methoxy-3-(2-nitro-phenylsulfanyl)-indol-1-ylmethyl]-pyridinium chloride (Compound 10)




20

To a solution of 6-Methoxy-3-(2-nitro-phenylsulfanyl)-1-pyridin-2-ylmethyl-1*H*-indole in methylene chloride was passed HCl-gas till the organic salt crystallised. The title compound was isolated by filtration.

- LC-MS : 4.04 min ($MH^+ = 392$); 1H NMR (DMSO): δ 3.78 (s, 3H, OCH₃), 5.78 (s, 2H, NCH₂), 6.77 (dd, 1H, $J_1 = 7.8$ Hz, $J_2 = 2.0$ Hz), 6.93 (dd, 1H, $J_1 = 7.8$ Hz, $J_2 = 0.8$ Hz), 7.22 (d, 1H, $J = 7.8$ Hz), 7.26 (d, 1H, $J = 2.0$ Hz), 7.34 (m, 2H), 7.48-7.52 (m, 1H), 7.59 (t, 1H, $J = 7.0$ Hz), 7.96 (s, 1H), 8.08 (t, 1H, $J = 7.0$ Hz), 8.27 (dd, 1H, $J_1 = 7.8$ Hz, $J_2 = 1.6$ Hz), 8.74 (d, 1H, $J = 6.7$ Hz).

10 Example 16

1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1*H*-indol-6-ylamine (Compound 57, Structure 23 of Scheme V, where $R^1 = 3,5$ -difluorophenyl, $R^2 = 2$ -nitrophenyl, $R^3 = R^4 = H$)



- 15 a) 1-(3,5-Difluoro-benzyl)-6-nitro-1*H*-indole (Structure 19 of Scheme V, where $R^1 = 3,5$ -difluorophenyl, $R^3 = R^4 = H$)

To a solution of 6-nitroindole (162 mg, 1.0 mmol) in DMF (4 mL) was added NaH (60% dispersion on oil; 80 mg, 2.0 mmol). The resulting dark solution was stirred at room temperature for 15 minutes. 3,5-difluorobenzyl bromide (129 μ L, 1.1 mmol) was added. The reaction mixture was stirred overnight, poured into acidified water and extracted twice with ethyl acetate. The combined organic layers were dried over Na₂SO₄ and concentrated *in vacuo*.

The product was purified over a SiO₂ column (heptane/ethyl acetate 9:1) to give the title compound as a yellow solid (270 mg, yield = 94%).

- 25 LCMS: 5.54 min (100.0%, $MH^+ = 289$); 1H NMR (CDCl₃): δ 5.40 (s, 2H, CH₂Ar), 6.58 (m, 2H), 6.71 (d, 1H, $J = 3.5$ Hz), 6.75 (m, 1H), 7.26 (s, 1H), 7.41 (d, 1H, $J = 3.5$ Hz), 7.71 (d, 1H, $J = 8.3$ Hz), 8.05 (dd, 1H, $J_1 = 8.3$ Hz, $J_2 = 2.0$ Hz), 8.21 (d, 1H, $J = 2.0$ Hz).

b) 1-(3,5-Difluoro-benzyl)-1*H*-indol-6-ylamine (Structure 20 of Scheme V, where $R^1 =$ 3,5-difluorophenyl, $R^3=R^4=H$)

To a solution of 1-(3,5-Difluoro-benzyl)-6-nitro-1*H*-indole (144 mg, 0.5 mmol) in ethanol (20 mL) was added hydrochloric acid 37% (80 μ L) and $SnCl_2 \cdot 2H_2O$ (600 mg). The reaction mixture was stirred at 60 °C for 40 h and was then concentrated *in vacuo*. The residue was poured into ethyl acetate and a concentrated $NaHCO_3$ -solution was added. The two-layer phase system was filtered over decalite to get rid of the tin salts and the filtrate was twice extracted with ethyl acetate. The combined organic layers were dried over Na_2SO_4 and concentrated *in vacuo*. The product was purified over a SiO_2 column (heptane/ethyl acetate 9:1) to give the title compound as a colourless oil (89 mg, yield = 69%).

LCMS: 3.20 min (97.7%, $MH^+ = 259$); 1H NMR ($CDCl_3$): δ 5.12 (s, 2H, CH_2Ar), 6.52 (m, 2H), 6.56 (d, 1H, $J = 3.5$ Hz), 6.63 (m, 1H), 7.97 (dd, 1H, $J_1 = 8.3$ Hz, $J_2 = 2.0$ Hz), 7.13 (d, 1H, $J = 3.5$ Hz), 7.17 (d, 1H, $J = 2.0$ Hz), 7.55 (d, 1H, $J = 8.3$ Hz).


c) 1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1*H*-indol-6-ylamine (Compound 57, Structure 23 of Scheme V, where $R^1 =$ 3,5-difluorophenyl, $R^2 =$ 2-nitrophenyl, $R^3=R^4=H$)

To a solution of 1-(3,5-Difluoro-benzyl)-1*H*-indol-6-ylamine (25 mg, 0.1 mmol) in DCM (4 mL) was added 2-nitrobenzenesulfenyl chloride. The resulting yellow solution was stirred at room temperature for 1 hour. The mixture was concentrated *in vacuo* and purified over a SiO_2 column (heptane/ethyl acetate 9:1) to give the title compound as an orange solid (11.6 mg, yield = 28%).

LCMS: 5.11 min (100.0%, $MH^+ = 412$); 1H NMR ($CDCl_3$): δ 5.22 (s, 2H, CH_2Ar), 6.56 (m, 2H), 6.65 (m, 1H), 6.84 (dd, 1H, $J_1 = 8.2$ Hz, $J_2 = 1.2$ Hz), 7.00 (dd, 1H, $J_1 = 8.2$ Hz, $J_2 = 1.2$ Hz), 7.19 (m, 1H), 7.30 (d, 1H, $J = 1.2$ Hz), 7.31 (s, 1H), 7.37 (d, 1H, $J = 8.2$ Hz), 7.49 (m, 1H), 8.27 (dd, 1H, $J_1 = 8.2$ Hz, $J_2 = 1.2$ Hz).

Example 17

N-[1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1*H*-indol-6-yl]-acetamide
 (Compound 58, Structure 22 of Scheme V, where $R^1 = 3,5$ -difluorophenyl, $R^2 = 2$ -nitrophenyl, $R^3 = R^4 = H$, $Z = \text{methyl}$)



General method 5: Reaction of an acid anhydride with a 6-aminoindole of structure 20 to give compounds of structure 21, followed by sulfanylation of the indole 3-position to give compounds of structure 22 (Scheme V).

a) *N*-[1-(3,5-Difluoro-benzyl)-1*H*-indol-6-yl]-acetamide (Structure 21 of Scheme V, where $R^1 = 3,5$ -difluorophenyl, $R^3 = R^4 = H$, $Z = \text{methyl}$)

To a solution of 1-(3,5-Difluoro-benzyl)-1*H*-indol-6-ylamine (25 mg, 0.1 mmol) in DCM (4 mL) was added pyridine (25 μ L) and acetic anhydride (9.8 μ L, 0.11 mmol) and stirred overnight at room temperature. The reaction mixture was poured into 10 mL of water and neutralised with NaHCO_3 and extracted twice with DCM. The combined organic layers were dried over Na_2SO_4 and concentrated *in vacuo* to give the crude compound as a off white solid (23.3 mg, yield = 84%). The product was used without further purification.

LCMS: 4.10 min (97.4%, $\text{MH}^+ = 301$); ^1H NMR (CDCl_3): δ 2.18 (s, 3H, CH_3CON), 5.28 (s, 2H, CH_2Ar), 6.53 (d, 1H, $J = 3.1$), 6.58 (m, 2H), 6.68 (m, 1H), 6.87 (dd, 1H, $J_1 = 8.3$ Hz, $J_2 = 1.6$ Hz), 7.07 (d, 1H, $J = 3.1$ Hz), 7.55 (d, 1H, $J = 8.3$ Hz), 7.90 (s, 1H).

b) *N*-[1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1*H*-indol-6-yl]-acetamide
 (Compound 58, Structure 22 of Scheme V, where $R^1 = 3,5$ -difluorophenyl, $R^2 = 2$ -nitrophenyl, $R^3 = R^4 = H$, $Z = \text{methyl}$)


To a solution of *N*-[1-(3,5-Difluoro-benzyl)-1*H*-indol-6-yl]-acetamide (23.3 mg, 0.08 mmol) in diethyl ether (4 mL) was added 2-nitrobenzenesulfonyl chloride (14.7 mg,

0.09 mmol). The resulting yellow solution was stirred overnight at room temperature. The mixture was concentrated *in vacuo* and purified over a SiO₂ column (heptane/ethyl acetate 9:1) to give the title compound as a yellow solid (35.0 mg, yield = 96%).

LCMS: 4.26 min (98.6%, MH⁺ = 454); ¹H NMR (DMSO): δ 2.03 (s, 3H, CH₃CON), 5.51 (s, 2H, CH₂Ar), 6.89 (dd, 1H, J = 8.3 Hz, J2 = 1.2 Hz), 6.95 (m, 2H), 7.17 (dd, 1H, J1 = 8.3 Hz, J2 = 2.0 Hz), 7.19 (m, 1H), 7.27 (d, 1H, J = 8.3 Hz), 7.34 (m, 1H), 7.49 (m, 1H), 7.97 (d, 1H, J = 1.2 Hz), 8.02 (s, 1H), 8.27 (dd, 1H, J1 = 8.3 Hz, J2 = 1.2 Hz).

Example 18

- 10 *N*-[1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1*H*-indol-6-yl]-2,2,2-trifluoro-acetamide (Compound 59, Structure 22 of Scheme V, where R¹ = 3,5-difluorophenyl, R² = 2-nitrophenyl, R³=R⁴=H, Z = trifluoromethyl)



According to General method 5, step a): *N*-[1-(3,5-Difluoro-benzyl)-1*H*-indol-6-yl]-2,2,2-trifluoro-acetamide was prepared using 1-(3,5-Difluoro-benzyl)-1*H*-indol-6-ylamine (17.7 mg, 69 μmol), pyridine (17 μL) and trifluoroacetic anhydride (9.68 μL, 70 μmol). The compound was purified over a SiO₂ column (heptane/ethyl acetate 9:1) to give the title compound as a white solid (10.9 mg, yield = 61%).

LCMS: 4.41 min (82.1%, MH⁺ = 355)

- 20 According to General method 5, step b): *N*-[1-(3,5-Difluoro-benzyl)-3-(2-nitro-phenylsulfanyl)-1*H*-indol-6-yl]-2,2,2-trifluoro-acetamide was prepared using *N*-[1-(3,5-Difluoro-benzyl)-1*H*-indol-6-yl]-2,2,2-trifluoro-acetamide (10.9 mg, 31 μmol) and 2-nitrobenzenesulfonylchloride (5.7 mg, 37 μmol). The compound was purified over an HPLC column (MeCN/H₂O) to give the title compound as a yellow solid (10.2 mg, yield = 65%).

LCMS: 4.65 min (89.0%, MH⁺ = 508); ¹H NMR (DMSO): δ 5.56 (s, 2H, CH₂Ar), 6.87 (dd, 1H, J = 8.3 Hz, J2 = 1.2 Hz), 7.01 (m, 2H), 7.20 (m, 1H), 7.35 (m, 1H), 7.37 (dd,

Cpd. number	X	R ¹	R ²	R ³	R ⁴	R ⁵	ago EC50	ago efficacy	ant EC50	ant efficacy
5	S	2-fluorophenyl	2-nitrophenyl	H	H	OMe	+	+	-	-
6	S	2-methyl,3-nitrophenyl	2-nitrophenyl	H	H	OMe	+	+	-	-
7	S	2-methylphenyl	2-nitrophenyl	H	H	OMe	++	++	-	-
8	S	2-nitrophenyl	2-nitrophenyl	H	H	OMe	++	+	-	-
9	S	2-pyridyl	2-nitrophenyl	H	H	OMe	+++	+	+++	±
10	S	2-pyridyl HCl salt	2-nitrophenyl	H	H	OMe	++	+	-	-
11	S	2-tetrahydropyrany	2-nitrophenyl	H	H	OMe	++	+	-	-
12	S	2-trifluoromethylphenyl	2-nitrophenyl	H	H	OMe	+	+	-	-
13	S	3-(5-methylisoxazolyl)	2-nitrophenyl	H	H	OMe	+	±	-	-
14	S	3,4-dichlorophenyl	2-nitrophenyl	H	H	OMe	+	+	-	-
15	S	3,5-dichlorophenyl	2-nitrophenyl	H	H	OMe	++	++	-	-
16	S	3,5-difluorophenyl	2-(hydroxymethylphenyl)	H	H	OMe	++	+	++	±
17	S	3,5-difluorophenyl	2-(N-methylbenzamide)	H	H	OMe	-	-	+	+
18	S	3,5-difluorophenyl	2-benzamide	H	H	OMe	++	+	++	±
19	S	3,5-difluorophenyl	2-benzoic acid methyl ester	H	H	OMe	+	+	-	-
20	S	3,5-difluorophenyl	2-cyanophenyl	H	H	OMe	++	++	-	-
21	S	3,5-difluorophenyl	2-methoxyphenyl	H	H	OMe	++			
22	S	3,5-difluorophenyl	2-nitrophenyl	H	H	Br	++	++	-	-
23	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CF ₃	+++	++	-	-
24	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CH ₂ OH	++	++	-	-
25	S	3,5-difluorophenyl	2-nitrophenyl	H	H	Cl	+++	++	-	-
26	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CN	+++	++	-	-
27	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CO(1-pyrrolidinyl)	++	+	-	-
28	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CO(4-morpholinyl)	++	+	-	-
29	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CO[1-(4- methylpiperazinyl)]	++	++	-	-
30	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CO ₂ H	+	+	-	-
31	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CO ₂ Me	+++	+	-	-
32	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONH ₂	++	+	+	±
33	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONH ₂	+++	++	-	-
34	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ (2-furanyl)	++	+	-	-
35	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ (3-pyridyl)	++	±	-	-
36	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ CH ₂ Nmc ₂	++	+	-	-

Cpd. number	X	R ¹	R ²	R ³	R ⁴	R ⁵	ago EC50	ago efficacy	ant EC50	ant efficacy
37	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ CH ₂ OH	+++	++	-	-
38	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ CH ₂ OMe	+++	++	-	-
39	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ CO ₂ Me	++	++	-	-
40	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ CONMe ₂	++	+	-	-
41	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ cPr	+++	+	-	-
42	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ iPr	++	+	-	-
43	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHCH ₂ Ph	++	±	+	±
44	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHcPr	+++	+	-	-
45	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHt	+++	++	-	-
46	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHiPr	++	+	++	±
47	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHMe	+++	+	-	-
48	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONHnPr	++	+	+	±
49	S	3,5-difluorophenyl	2-nitrophenyl	H	H	CONMe ₂	+++	+	-	-
50	S	3,5-difluorophenyl	2-nitrophenyl	H	H	F	++	++	-	-
51	S	3,5-difluorophenyl	2-nitrophenyl	F	H	H	+++	++	-	-
52	S	3,5-difluorophenyl	2-nitrophenyl	H	H	H	++	++	-	-
53	S	3,5-difluorophenyl	2-nitrophenyl	H	F	H	++	++	-	-
54	S	3,5-difluorophenyl	2-nitrophenyl	Cl	H	H	++	++	-	-
55	S	3,5-difluorophenyl	2-nitrophenyl	Me	H	H	++	+	-	-
56	S	3,5-difluorophenyl	2-nitrophenyl	H	OH	H	+++			
57	S	3,5-difluorophenyl	2-nitrophenyl	H	H	NH ₂	+	+	-	-
58	S	3,5-difluorophenyl	2-nitrophenyl	F	H	NHAc	+++	++	-	-
59	S	3,5-difluorophenyl	2-nitrophenyl	H	H	NHCOCF ₃	+++	+	-	-
60	S	3,5-difluorophenyl	2-nitrophenyl	H	H	NO ₂	+++	++	-	-
61	S	3,5-difluorophenyl	2-nitrophenyl	H	H	OH	+++			
62	SO	3,5-difluorophenyl	2-nitrophenyl	H	H	OMe	+++	++	-	-
2										
63	S	3,5-difluorophenyl	2-nitrophenyl	H	H	OMc	+++	++	-	-
64	SO	3,5-difluorophenyl	2-nitrophenyl	H	H	OMe	-	±	+	+
65	S	3,5-difluorophenyl	2-pyridyl	H	H	OMc	++	++	-	-
66	S	3,5-difluorophenyl	2-pyridyl-N-oxide	H	H	OMe	+	+	++	±
67	S	3-chlorophenyl	2-nitrophenyl	H	H	OMe	++	++	-	-
68	S	3-cyanophenyl	2-nitrophenyl	H	H	OMc	+++	++	-	-
69	S	3-fluoro,5-	2-nitrophenyl	H	H	OMe	++	++	-	-


Cpd. number	X	R ¹	R ²	R ³	R ⁴	R ⁵	ago EC50	ago efficacy	ant EC50	ant efficacy
70	S	trifluoromethylphenyl 3-fluoro,6-	2-nitrophenyl	H	H	OMe	++	+	-	-
71	S	trifluoromethylphenyl 3-fluorophenyl	2-nitrophenyl	H	H	OMe	+++	++	-	-
72	S	3-methoxyphenyl	2-nitrophenyl	H	H	OMe	++	+	-	-
73	S	3-methylphenyl	2-nitrophenyl	H	H	OMe	++	++	-	-
74	S	3-nitrophenyl	2-nitrophenyl	H	H	OMe	+++	++	-	-
75	S	3-pyridyl	2-nitrophenyl	H	H	OMe	+++	++	-	-
76	S	3-trifluoromethyl,4- chlorophenyl	2-nitrophenyl	H	H	OMe	+	+	-	-
77	S	3- trifluoromethyloxyphenyl	2-nitrophenyl	H	H	OMe	++	++	-	-
78	S	3-trifluoromethylphenyl	2-nitrophenyl	H	H	OMe	++	++	-	-
79	S	4-(2-methylthiazolyl)	2-nitrophenyl	H	H	OMe	++	+	-	-
80	S	4-(3,5- dimethylisoxazolyl)	2-nitrophenyl	H	H	OMe	+	±	+	±
81	S	4-chlorophenyl	2-nitrophenyl	H	H	OMe	+	+	+++	+
82	S	4-cyanophenyl	2-nitrophenyl	H	H	OMe	++	+	-	-
83	S	4-fluorophenyl	2-nitrophenyl	H	H	OMe	++	++	-	-
84	S	4-methoxyphenyl	2-nitrophenyl	H	H	OMe	+	±	±	++
85	S	4-methylphenyl	2-nitrophenyl	H	H	OMe	+	+	-	±
86	S	4-morpholinyl	2-nitrophenyl	H	H	OMe	±	±	+	+
87	S	5-(2-chlorothiazolyl)	2-nitrophenyl	H	H	OMe	++	+	-	-
88	S	5-(2-chlorothiophenyl)	2-nitrophenyl	H	H	OMe	++	++	-	-
89	S	Cyclohexyl	2-nitrophenyl	H	H	OMe	++	+	+++	±
90	S	Phenyl	2-nitrophenyl	H	H	H	+	+	-	-
91	S	Phenyl	2-nitrophenyl	H	OM	H	-	-	±	++
92	S	Phenyl	2-nitrophenyl	H	H	OMe	+++	++	-	-

note 1: EC50 scale: +++ <5 nM; ++ between 5 and 100 nM; + <1000 nM; ± between 1000 and 10000 nM; - >10000 nM

note 2: efficacy scale (maximal intrinsic activity, i.e. intrinsic activity observed in the presence of 100 nM DHT, was set at 1.00): ++ ≥ 0.80 ; + between 0.50 and 0.80; \pm between 0.20 and 0.50; - < 0.20

Claims

1. A compound having the formula



5 wherein

X is S, SO or SO₂;

R¹ is a 5-6-membered monocyclic, hetero- or homocyclic, saturated or unsaturated ring structure optionally substituted with one or more substituents selected from the group consisting of halogen, CN, (1C-4C)fluoroalkyl, nitro, (1C-4C)alkyl, (1C-4C)alkyloxy or (1C-4C)fluoroalkoxy;

R² is 2-nitrophenyl, 2-cyanophenyl, 2-hydroxymethyl-phenyl, 2-pyridyl, 2-pyridyl-N-oxide, 2-benzamide, 2-benzoic acid methyl ester or 2-methoxyphenyl;

R³ is H, halogen or (1C-4C)alkyl;

R⁴ is H, OH or halogen;

R⁵ is H, OH, (1C-4C)alkoxy, NH₂, CN, halogen, (1C-4C)fluoroalkyl, NO₂, hydroxy(1C-4C)alkyl, CO₂H or

R⁴ is NHR⁶, wherein R⁶ is (1C-6C)acyl optionally substituted with one or more halogens, or

R⁵ is C(O)N(R⁸, R⁹), wherein R⁸ and R⁹ each independently are H, (1C-6C)cycloalkyl, or

R⁸ and R⁹ form together with the N a heterocyclic 5-6-membered saturated or unsaturated ring optionally substituted with (1C-4C)alkyl, or

R⁸ and R⁹ each independently are CH₂R¹⁰, wherein R¹⁰ is H, (1C-5C)alkyl, hydroxy(1C-3C)alkyl, (1C-4C)alkylester of carboxy(1C-4C)alkyl, (1C-3C)alkoxy(1C-3C)alkyl, (mono- or di(1C-4C)alkyl)amino, (mono- or di(1C-4C)alkyl)aminocarbonyl, or a 5-6-membered monocyclic, homo- or heterocyclic, aromatic or non-aromatic ring;

or a salt or hydrate form thereof.

2. A compound according to claim 1, characterised in that

R^1 is a 5-6-membered monocyclic, hetero- or homocyclic, saturated or unsaturated ring structure optionally substituted with one or more substituents selected from the group consisting of halogen, CN, CF_3 , nitro, methoxy, trifluoromethoxy or methyl;

5 R^2 is 2-nitrophenyl, 2-cyanophenyl, 2-hydroxymethyl-phenyl, 2-pyridyl, 2-pyridyl-N-oxide, benzamide, 2-benzoic acid methyl ester or 2-methoxyphenyl;

R^3 is H, halogen or (1C-2C)alkyl;

R^4 is H or F.

10 3. A compound according to claim 2, characterised in that

R^5 is H, (1C-4C)alkoxy, CN, Hal, (1C-4C)fluoroalkyl, NO_2 , or

R^5 is NHR^6 , wherein R^6 is (1C-6C)acyl optionally substituted with one or more halogens, or

15 R^5 is $C(O)N(R^8, R^9)$, wherein R^8 and R^9 each independently are H, (1C-6C)cycloalkyl, or

R^8 and R^9 form together with the N a heterocyclic 5-6-membered saturated or unsaturated ring optionally substituted with (1C-4C)alkyl, or

20 R^8 and R^9 each independently are CH_2R^{10} , wherein R^{10} is H, (1C-5C)alkyl, hydroxy(1C-3C)alkyl, carboxy(1C-4C)alkyl, (1C-3C)alkoxy(1C-3C)alkyl, (mono- or di(1C-4C)alkyl)amino, (mono- or di(1C-4C)alkyl)-aminocarbonyl, or a 5-6-membered monocyclic, homo- or heterocyclic, aromatic or non-aromatic ring.

25 4. A compound according to claim 3, characterised in that

R^3 is H or halogen;

R^4 is H;

R^5 is H, OH, (1C-4C)alkyloxy, CN, F, Cl, CF_3 , NO_2 , or

30 R^5 is NHR^6 , wherein R^6 is (1C-3C)acyl optionally substituted with one or more halogens or

R^5 is $C(O)N(R^8, R^9)$, wherein R^8 and R^9 each independently are H, (1C-4C)cycloalkyl, or

35 R^8 and R^9 each independently are CH_2R^{10} , wherein R^{10} is H, (1C-5C)alkyl, hydroxy(1C-3C)alkyl, (1C-2C)alkylester of carboxy(1C-2C)alkyl, (1C-3C)alkyloxy(1C-3C)alkyl, (1C-4C)cycloalkyl, (mono- or di(1C-

4C)alkyl)amino, (mono- or di(1C-4C)alkyl)aminocarbonyl, or a 5-membered heterocyclic ring.

5. A compound according to claim 4, characterised in that

5 X is S or SO₂.

R² is 2-nitrophenyl, 2-hydroxymethyl-phenyl, 2-benzamide, 2-methoxyphenyl, 2-cyanophenyl or 2-pyridyl;

R³ is H or F;

R⁵ is H, OH, (1C-2C)alkyloxy, CN, F, Cl, CF₃, NO₂, or

10 R⁵ is NHR⁶, wherein R⁶ is acetyl or trifluoroacetyl, or


R⁵ is C(O)N(R⁸, R⁹), wherein R⁸ is H and R⁹ is H, cyclopropyl or

R⁹ is CH₂R¹⁰, wherein R¹⁰ is H, (1C-2C)alkyl, hydroxy(1C-2C)alkyl, methoxy(1C-2C)alkyl, cyclopropyl.

Abstract

This invention provides non-steroidal compounds with affinity for the androgen receptor and utility for androgen-receptor related treatments, having a structure

5 according to the formula



wherein X is S, SO or SO₂; R¹ is a 5-6-membered monocyclic, hetero- or homocyclic, saturated or unsaturated ring structure optionally substituted with one or more substituents selected from the group consisting of halogen, CN, (1C-4C)fluoroalkyl, nitro, (1C-4C)alkyl, (1C-4C)alkyloxy or (1C-4C)fluoroalkoxy; R² is 2-nitrophenyl, 2-cyanophenyl, 2-hydroxymethyl-phenyl, 2-pyridyl, 2-pyridyl-N-oxide, 2-benzamide, 2-benzoic acid methyl ester or 2-methoxyphenyl; R³ is H, halogen or (1C-4C)alkyl; R⁴ is H, OH or halogen; R⁵ is H, OH, (1C-4C)alkoxy, NH₂, CN, halogen, (1C-4C)fluoroalkyl, NO₂, hydroxy(1C-4C)alkyl, CO₂H, NHR⁶ or C(O)N(R⁸,R⁹). Herein is R⁶ (1C-6C)acyl optionally substituted with one or more halogens, and are R⁸ and R⁹ each independently H, (1C-6C)cycloalkyl, CH₂R¹⁰ or form together with the N a heterocyclic 5-6-membered saturated or unsaturated ring optionally substituted with (1C-4C)alkyl, where by R¹⁰ is H, (1C-5C)alkyl, hydroxy(1C-3C)alkyl, (1C-4C)alkylester of carboxy(1C-4C)alkyl, (1C-3C)alkoxy(1C-3C)alkyl, (mono- or di(1C-4C)alkyl)amino, (mono- or di(1C-4C)alkyl)aminocarbonyl, or a 5-6-membered monocyclic, homo- or heterocyclic, aromatic or non-aromatic ring.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.